

The ALSFRS-R-SE is an assessment of symptoms and motor deficits commonly associated with ALS that patients can complete themselves or with the help of others, such as a family member, caregiver, or healthcare professional, based on an interview with the patient.

Please carefully read the answer options and explanations. Then, select an option that reflects the current symptoms and motor functions compared to the situation before the onset of ALS.

1. Speech

Please choose one of the following answers:

- 4 Normal
Speech is normal.
- 3 Noticeable speech disturbance
Speech is slightly slurred or voice is rough, noticed by oneself or someone else.
- 2 Repetition is needed for understanding
Speech is moderately slurred; sometimes words or parts of sentences need to be repeated to be understood.
- 1 Speech combined with non-verbal communication
Speech is severely slurred; writing things down or use of communication aids are often needed to communicate.
- 0 Loss of understandable speech
Speech is lost; writing things down, communication devices, or other supportive methods are always needed.

2. Salivation

Please choose one of the following answers:

- 4 Normal
No excess saliva is accumulated in the mouth.
- 3 Slight but definite excess saliva in the mouth. May have night time drooling
More saliva is kept in the mouth. Saliva may leak out at night, but not during the day.
- 2 Moderate excess saliva. May have minimal day time drooling
A cloth or tissue is sometimes used to wipe the corners of the mouth.
- 1 Marked excess saliva with some drooling
A cloth or tissue is often used, but not always.
- 0 Severe excess saliva with marked drooling
A cloth or suction device is always required.

continued on the next page

3. Swallowing

Please choose one of the following answers:

- 4 Normal
Swallowing is normal.
- 3 Minor swallowing difficulties and occasional choking
Eating takes more time, or smaller bites are taken and swallowed carefully. Occasional choking or more frequent coughing may occur.
- 2 Moderate swallowing difficulties and dietary consistency changes
The texture of food and drinks is changed, or certain foods (e.g., meat, dry biscuits, nuts) are avoided. High-calorie drinks or thickening powders are used.
- 1 Severe swallowing difficulties. Supplementary tube feeding
A feeding tube has been placed or recommended to maintain weight. Some food or drink can be taken by mouth.
- 0 Unable to swallow. Exclusive tube feeding
A feeding tube has been placed. Eating or drinking by mouth is not possible.

4. Handwriting

Comment: The subject of this assessment is writing with one's dominant hand (writing hand) in the usual posture.

Please choose one of the following answers:

- 4 Normal
Writing is normal.
- 3 Slow or sloppy but all words are legible
Writing is slower and less clear, but the words can be written.
- 2 Not all words are legible
Some words are not legible. Tools may be used to make writing clearer.
- 1 No words are legible, but can still grip a pen
A pen can be held. Only the signature or one's own name, but no other words can be written.
- 0 Cannot grip a pen
A pen cannot be held.

5.a Cutting food and handling utensils

Comment: For persons who do not use a feeding tube.

Please choose one of the following answers:

- 4 Normal
Food is handled without difficulty using standard utensils such as a knife, spoon, fork, or chopsticks.
- 3 Somewhat slow and clumsy, but no help needed
Food is handled more slowly or clumsily with standard utensils, but without help.
- 2 Can cut most foods, although slowly and clumsily. Some help needed
Food can be handled independently, sometimes using special utensils such as adapted knives or forks. Help is occasionally needed to cut certain foods.
- 1 Food must be cut by someone else. Can eat slowly on their own
Food can be handled independently, if needed using special utensils. Help is always needed to cut solid foods.
- 0 Cannot eat on their own. Always help needed
Food cannot be handled independently. Help is always needed.

continued on the next page

5.b Handling of feeding tube

Comment: For persons who use a feeding tube.

Please choose one of the following answers:

- 4 Normal
The feeding tube can be handled without difficulties.
- 3 Somewhat slow and clumsy, but handling of feeding tube can be performed independently
The feeding tube is handled slowly or clumsily, but without help.
- 2 Some help is needed when handling closures and lids
The feeding tube is handled independently, bur help is needed with opening or closing some parts, such as caps or lids.
- 1 Minimal assistance to care giver in the handling of feeding tube
The feeding tube cannot be handled independently. Help is mostly needed, and only small parts can be done alone.
- 0 Unable to perform any aspects of the task
The feeding tube cannot be handled independently. Help is always needed.

6. Dressing and hygiene

Please choose one of the following answers:

- 4 Normal
Dressing and hygiene are done without difficulties.
- 3 Independent and complete self-care with effort or decreased efficiency
Dressing and hygiene take longer but can be done without help.
- 2 Intermittent help or substitute methods
Dressing and hygiene are done with substitute methods, such as easy clothing or sitting while dressing. Help is sometimes needed.
- 1 Needs help with self-care
Dressing and hygiene cannot be done independently. Help is mostly needed, and only small parts can be done alone.
- 0 Total dependence
Dressing and hygiene cannot be done independently. Help is always needed.

7. Turning in bed and adjusting bed clothes

Please choose one of the following answers:

- 4 Normal
Turning in bed and handling blankets is normal.
- 3 Somewhat slow and clumsy, but no help is needed
Turning over in bed and handling the blankets are slower and clumsy, but without help.
- 2 Can turn alone or adjust sheets, but with great difficulty
Turning over in bed and handling the blankets can be done with great effort. Either action sometimes needs help.
- 1 Can initiate, but not turn or adjust sheets alone
Turning over in bed or handling the blankets can be started, but help is needed to complete them.
- 0 Total dependence
Turning over in bed and handling the blankets cannot be done independently. Help is always needed.

continued on the next page

8. Walking

Please choose one of the following answers:

- 4 Normal
Walking is normal.
- 3 Minor walking difficulties
Walking is slower, feels steady, or leads to occasional tripping. No assistance is needed when outside.
- 2 Walking with assistance
Walking needs assistance, regularly when outside, either by holding onto support by using a foot lifter or walking aid, or with help from someone else.
- 1 Unable to walk. Non-ambulatory leg movement
Walking is not possible, even with walking aids or the help of someone else. Standing is possible with assistance, for example, when transferring from bed to chair. The legs can be moved purposefully.
- 0 Unable to walk. No purposeful leg movement
Standing is not possible, even with assistance. The legs cannot be moved purposefully.

9. Climbing the stairs

Please choose one of the following answers:

- 4 Normal
Climbing the stairs is normal.
- 3 Slow
Climbing the stairs is slowed but steady.
- 2 Mild unsteadiness or fatigue
Climbing the stairs is slowed and unsteady, and rest may be needed. Assistance, such as holding a handrail, is not regularly required.
- 1 Needs assistance
Climbing the stairs requires assistance, such as a handrail or help from someone else.
- 0 Cannot climb
Climbing the stairs is not possible, even with help.

10. Shortness of breath

Please choose one of the following answers:

- 4 No breathing difficulty
No shortness of breath is felt when doing daily activities.
- 3 Occurs when walking
Shortness of breath is felt when walking at a normal speed or doing everyday tasks that need some effort.
- 2 Occurs with one or more of the following: eating, bathing, dressing, prolonged speaking
Shortness of breath is felt during light activities.
- 1 Difficulty breathing when at rest, including either sitting or lying down
Shortness of breath is felt when resting, such as sitting or lying down.
- 0 Significant difficulty breathing, mechanical respiratory support may be needed
Intense shortness of breath is felt when resting. Ventilation support, such as a breathing mask or tracheostomy, is needed.

continued on the next page

11. Sleep disturbance due to shortness of breath

Comment: If a breathing mask is used at night but sleep is also possible without it, sleep should be rated for nights without the mask.

Please choose one of the following answers:

- 4 None
Sleep is not disturbed due to shortness of breath.
- 3 Some difficulty sleeping at night due to shortness of breath. More than two pillows are not routinely used
Sleep is disturbed due to shortness of breath. Sleeping on the side improves the breathing.
- 2 Regular use of more than two pillows required for sleeping
Sleep is difficult due to shortness of breath. Sleeping with three or more pillows or by raising the head of the bed improves the breathing.
- 1 Can only sleep sitting up
Sleep is possible only when sitting up, either in bed or in a chair.
- 0 Unable to sleep
Sleep is possible only with ventilation support such as breathing mask or tracheostomy.



12. Ventilation support, such as a breathing mask or tracheostomy

Please choose one of the following answers:

- 4 None
No ventilation support is used. An existing treatment of sleep apnoea with a CPAP mask is not counted as ventilation support.
- 3 Intermittent use of breathing mask
A breathing mask is used not every night or not throughout the entire night.
- 2 Continuous use of breathing mask during the night
A breathing mask is used every night and sometimes during the day, for a total of 8 to 22 hours.
- 1 Continuous use of non-invasive ventilation during the night and day
A breathing mask is used nearly all night and day, for more than 22 hours.
- 0 Invasive ventilation with tracheostomy
Invasive ventilation is used through a tracheostomy.

